

OSCILLATION CRITERIA FOR FUNCTIONAL DIFFERENTIAL EQUATIONS

by

I.P. Stavroulakis
Department of Mathematics
University of Ioannina
451 10 Ioannina, Greece
ipstav@cc.uoi.gr

ABSTRACT

Consider the first-order linear delay differential equation

$$x'(t) + p(t)x(\tau(t)) = 0, \quad t \geq t_0, \quad (1)$$

and the second-order linear delay equation

$$x''(t) + p(t)x(\tau(t)) = 0, \quad t \geq t_0, \quad (2)$$

where $p \in C([t_0, \infty), \mathbb{R}^+)$, $\tau \in C([t_0, \infty), \mathbb{R})$, $\tau(t)$ is non-decreasing, $\tau(t) \leq t$ for $t \geq t_0$ and $\lim_{t \rightarrow \infty} \tau(t) = \infty$.

The most interesting oscillation criteria for Eq.(1), especially in the case where

$$0 < \liminf_{t \rightarrow \infty} \int_{\tau(t)}^t p(s)ds \leq \frac{1}{e} \quad \text{and} \quad \limsup_{t \rightarrow \infty} \int_{\tau(t)}^t p(s)ds < 1,$$

and for Eq. (2) when

$$\liminf_{t \rightarrow \infty} \int_{\tau(t)}^t \tau(s)p(s)ds \leq \frac{1}{e} \quad \text{and} \quad \limsup_{t \rightarrow \infty} \int_{\tau(t)}^t \tau(s)p(s)ds < 1$$

are presented.

⁰Key Words: Oscillation; delay differential equations.

2000 Mathematics Subject Classification: Primary 34K11; Secondary 34C10.

1 Introduction

The problem of establishing sufficient conditions for the oscillation of all solutions to the first-order differential equation

$$x'(t) + p(t)x(\tau(t)) = 0, \quad t \geq t_0, \quad (1)$$

and the second-order equation

$$x''(t) + p(t)x(\tau(t)) = 0, \quad t \geq t_0, \quad (2)$$

where $p \in C([t_0, \infty), \mathbb{R}^+)$ (here $\mathbb{R}^+ = [0, \infty)$), $\tau \in C([t_0, \infty), \mathbb{R})$, $\tau(t)$ is non-decreasing, $\tau(t) \leq t$ for $t \geq t_0$ and $\lim_{t \rightarrow \infty} \tau(t) = \infty$, has been the subject of many investigations. See, for example, [1-65] and the references cited therein.

By a solution of Eq.(1) [resp. Eq.(2)] we understand a continuously differentiable function defined on $[\tau(T_0), \infty)$ for some $T_0 \geq t_0$ and such that Eq.(1) [resp. Eq.(2)] is satisfied for $t \geq T_0$. Such a solution is called *oscillatory* if it has arbitrarily large zeros, and otherwise it is called *nonoscillatory*.

In this paper our main purpose is to present the state of the art on the oscillation of all solutions to Eq.(1) especially in the case where

$$0 < \liminf_{t \rightarrow \infty} \int_{\tau(t)}^t p(s) ds \leq \frac{1}{e} \quad \text{and} \quad \limsup_{t \rightarrow \infty} \int_{\tau(t)}^t p(s) ds < 1,$$

and for Eq.(2) when

$$\liminf_{t \rightarrow \infty} \int_{\tau(t)}^t \tau(s) p(s) ds \leq \frac{1}{e} \quad \text{and} \quad \limsup_{t \rightarrow \infty} \int_{\tau(t)}^t \tau(s) p(s) ds < 1.$$

2 Oscillation Criteria for Eq. (1)

In this section we study the delay equation

$$x'(t) + p(t)x(\tau(t)) = 0, \quad t \geq t_0. \quad (1)$$

The first systematic study for the oscillation of all solutions to Eq.(1) was made by Myshkis. In 1950 [42] he proved that every solution of Eq.(1) oscillates if

$$\limsup_{t \rightarrow \infty} [t - \tau(t)] < \infty \quad \text{and} \quad \liminf_{t \rightarrow \infty} [t - \tau(t)] \liminf_{t \rightarrow \infty} p(t) > \frac{1}{e}. \quad (C_1)$$

In 1972, Ladas, Lakshmikantham and Papadakis [33] proved that the same conclusion holds if

$$A := \limsup_{t \rightarrow \infty} \int_{\tau(t)}^t p(s) ds > 1. \quad (C_2)$$

In 1979, Ladas [32] established integral conditions for the oscillation of Eq.(1) with constant delay. Tomaras [54-56] extended this result to Eq.(1) with variable delay. For related results see Ladde [36-38]. The following most general result is due to Koplatadze and Canturija [25].

If

$$\alpha := \liminf_{t \rightarrow \infty} \int_{\tau(t)}^t p(s) ds > \frac{1}{e}, \quad (C_3)$$

then all solutions of Eq.(1) oscillate; If

$$\limsup_{t \rightarrow \infty} \int_{\tau(t)}^t p(s) ds < \frac{1}{e}, \quad (N_1)$$

then Eq.(1) has a nonoscillatory solution.

In 1982 Ladas, Sficas and Stavroulakis [35] and in 1984 Fukagai and Kusano [13] established oscillation criteria (of the type of conditions (C_2) and (C_3)) for Eq. (1) with *oscillating* coefficient $p(t)$.

It is obvious that there is a gap between the conditions (C_2) and (C_3) when the limit $\lim_{t \rightarrow \infty} \int_{\tau(t)}^t p(s) ds$ does not exist. How to fill this gap is an interesting problem which has been recently investigated by several authors.

In 1988, Erbe and Zhang [12] developed new oscillation criteria by employing the upper bound of the ratio $x(\tau(t))/x(t)$ for possible nonoscillatory solutions $x(t)$ of Eq.(1). Their result says that all the solutions of Eq.(1) are oscillatory, if $0 < \alpha \leq \frac{1}{e}$ and

$$A > 1 - \frac{\alpha^2}{4}. \quad (C_4)$$

Since then several authors tried to obtain better results by improving the upper bound for $x(\tau(t))/x(t)$.

In 1991, Jian [20] derived the condition

$$A > 1 - \frac{\alpha^2}{2(1-\alpha)}, \quad (C_5)$$

while in 1992, Yu and Wang [63] and Yu, Wang, Zhang and Qian [64] obtained the condition

$$A > 1 - \frac{1 - \alpha - \sqrt{1 - 2\alpha - \alpha^2}}{2}. \quad (C_6)$$

In 1990, Elbert and Stavroulakis [8] and in 1991 Kwong [30], using different techniques, improved (C_4) , in the case where $0 < \alpha \leq \frac{1}{e}$, to the conditions

$$A > 1 - \left(1 - \frac{1}{\sqrt{\lambda_1}}\right)^2 \quad (C_7)$$

and

$$A > \frac{\ln \lambda_1 + 1}{\lambda_1}, \quad (C_8)$$

respectively, where λ_1 is the smaller real root of the equation $\lambda = e^{\alpha\lambda}$.

In 1994, Koplatadze and Kvinikadze [26] improved (C_6) , while in 1998, Philos and Sficas [45] and in 1999, Zhou and Yu [65] and Jaroš and Stavroulakis [19] derived the conditions

$$A > 1 - \frac{\alpha^2}{2(1-\alpha)} - \frac{\alpha^2}{2}\lambda_1, \quad (C_9)$$

$$A > 1 - \frac{1-\alpha-\sqrt{1-2\alpha-\alpha^2}}{2} - \left(1 - \frac{1}{\sqrt{\lambda_1}}\right)^2, \quad (C_{10})$$

and

$$A > \frac{\ln \lambda_1 + 1}{\lambda_1} - \frac{1-\alpha-\sqrt{1-2\alpha-\alpha^2}}{2}, \quad (C_{11})$$

respectively.

Consider Eq.(1) and assume that $\tau(t)$ is continuously differentiable and that there exists $\theta > 0$ such that $p(\tau(t))\tau'(t) \geq \theta p(t)$ eventually for all t . Under this additional condition, in 2000, Kon, Sficas and Stavroulakis [22] and in 2003, Sficas and Stavroulakis [46] established the conditions

$$A > \frac{\ln \lambda_1 + 1}{\lambda_1} - \frac{1-\alpha-\sqrt{(1-\alpha)^2-4\Theta}}{2} \quad (2.1)$$

and

$$A > \frac{\ln \lambda_1}{\lambda_1} - \frac{1+\sqrt{1+2\theta-2\theta\lambda_1 M}}{\theta\lambda_1} \quad (2.2)$$

respectively, where

$$\Theta = \frac{e^{\lambda_1\theta\alpha} - \lambda_1\theta\alpha - 1}{(\lambda_1\theta)^2}$$

and

$$M = \frac{1-\alpha-\sqrt{(1-\alpha)^2-4\Theta}}{2}.$$

Remark 2.1 ([22], [46]) Observe that when $\theta = 1$, then $\Theta = \frac{\lambda_1 - \lambda_1 \alpha - 1}{\lambda_1^2}$, and (2.1) reduces to

$$A > 2\alpha + \frac{2}{\lambda_1} - 1, \quad (C_{12})$$

while in this case it follows that $M = 1 - \alpha - \frac{1}{\lambda_1}$ and (2.2) reduces to

$$A > \frac{\ln \lambda_1 - 1 + \sqrt{5 - 2\lambda_1 + 2\alpha\lambda_1}}{\lambda_1}. \quad (C_{13})$$

In the case where $\alpha = \frac{1}{e}$, then $\lambda_1 = e$, and (C₁₃) leads to

$$A > \frac{\sqrt{7 - 2e}}{e} \approx 0.459987065.$$

It is to be noted that as $\alpha \rightarrow 0$, then all the previous conditions (C₄) – (C₁₂) reduce to the condition (C₂), i.e.

$$A > 1.$$

However, the condition (C₁₃) leads to

$$A > \sqrt{3} - 1 \approx 0.732,$$

which is an essential improvement. Moreover (C₁₃) improves all the above conditions when $0 < \alpha \leq \frac{1}{e}$ as well. Note that the value of the lower bound on A can not be less than

$$\frac{1}{e} \approx 0.367879441.$$

Thus the aim is to establish a condition which leads to a value *as close as possible* to $\frac{1}{e}$. For illustrative purpose, we give the values of the lower bound on A under these conditions when $\alpha = \frac{1}{e}$.

(C ₄):	0.966166179
(C ₅):	0.892951367
(C ₆):	0.863457014
(C ₇):	0.845181878
(C ₈):	0.735758882
(C ₉):	0.709011646
(C ₁₀):	0.708638892
(C ₁₁):	0.599215896
(C ₁₂):	0.471517764
(C ₁₃):	0.459987065

We see that the condition (C_{13}) essentially improves all the known results in the literature.

Example 2.1 ([46]) Consider the delay differential equation

$$x'(t) + px(t - q \sin^2 \sqrt{t} - \frac{1}{pe}) = 0,$$

where $p > 0$, $q > 0$ and $pq = 0.46 - \frac{1}{e}$. Then

$$\alpha = \liminf_{t \rightarrow \infty} \int_{\tau(t)}^t p ds = \liminf_{t \rightarrow \infty} p(q \sin^2 \sqrt{t} + \frac{1}{pe}) = \frac{1}{e}$$

and

$$A = \limsup_{t \rightarrow \infty} \int_{\tau(t)}^t p ds = \limsup_{t \rightarrow \infty} p(q \sin^2 \sqrt{t} + \frac{1}{pe}) = pq + \frac{1}{e} = 0.46.$$

Thus, according to Remark 2.1, all solutions of this equation oscillate. Observe that none of the conditions (C_4) - (C_{12}) apply to this equation.

Following this historical (and chronological) review we also mention that in the case where

$$\int_{\tau(t)}^t p(s) ds \geq \frac{1}{e} \quad \text{and} \quad \lim_{t \rightarrow \infty} \int_{\tau(t)}^t p(s) ds = \frac{1}{e}$$

this problem has been studied in 1995, by Elbert and Stavroulakis [9], by Koza-kiewicz [28], Li [40,41] and in 1996, by Domshlak and Stavroulakis [6].

3 Oscillation Criteria for Eq. (2)

In this section we study the second-order delay equation

$$x''(t) + p(t)x(\tau(t)) = 0, \quad t \geq t_0, \quad (2)$$

For the case of ordinary differential equations, i.e. when $\tau(t) \equiv t$, the history of the problem began as early as in 1836 by the work of Sturm [47] and was continued in 1893 by A. Kneser [21]. Essential contribution to the subject was made by E. Hille, A. Wintner, Ph. Hartman, W. Leighton, Z. Nehari, and others (see the monograph by C. Swanson [48] and the references cited therein). In particular, in 1948 E. Hille [17] obtained the following well-known oscillation criteria. Let

$$\limsup_{t \rightarrow \infty} t \int_t^{+\infty} p(s) ds > 1 \quad (3.1)$$

or

$$\liminf_{t \rightarrow \infty} t \int_t^{+\infty} p(s) ds > \frac{1}{4}, \quad (3.2)$$

the conditions being assumed to be satisfied if the integral diverges. Then Eq.(2) with $\tau(t) \equiv t$ is oscillatory.

For the delay differential equation (2) earlier oscillation results can be found in the monographs by A. Myshkis [43] and S. Norkin [44]. In 1968 P. Waltman [57] and in 1970 J. Bradley [1] proved that (2) is oscillatory if

$$\int^{+\infty} p(t) dt = +\infty.$$

Proceeding in the direction of generalization of Hille's criteria, in 1971 J. Wong [60] showed that if $\tau(t) \geq \alpha t$ for $t \geq 0$ with $0 < \alpha \leq 1$, then the condition

$$\liminf_{t \rightarrow \infty} t \int_t^{+\infty} p(s) ds > \frac{1}{4\alpha} \quad (3.3)$$

is sufficient for the oscillation of Eq.(2). In 1973 L. Erbe [10] generalized this condition to

$$\liminf_{t \rightarrow \infty} t \int_t^{+\infty} \frac{\tau(s)}{s} p(s) ds > \frac{1}{4} \quad (3.4)$$

without any additional restriction on τ . In 1987 J. Yan [61] obtained some general criteria improving the previous ones.

An oscillation criterion of different type is given in 1986 by R. Koplatadze [23] and in 1988 by J. Wei [59], where it is proved that Eq.(2) is oscillatory if

$$\limsup_{t \rightarrow \infty} \int_{\tau(t)}^t \tau(s) p(s) ds > 1 \quad (C_2)'$$

or

$$\liminf_{t \rightarrow \infty} \int_{\tau(t)}^t \tau(s) p(s) ds > \frac{1}{e}. \quad (C_3)'$$

The conditions $(C_2)'$ and $(C_3)'$ are analogous to the oscillation conditions

$$A = \limsup_{t \rightarrow \infty} \int_{\tau(t)}^t p(s) ds > 1, \quad (C_2)$$

$$\alpha = \liminf_{t \rightarrow \infty} \int_{\tau(t)}^t p(s) ds > \frac{1}{e} \quad (C_3)$$

respectively, for the first order delay equation

$$x'(t) + p(t)x(\tau(t)) = 0. \quad (1)$$

The essential difference between (3.3),(3.4) and $(C_2)'$, $(C_3)'$ is that the first two can guarantee oscillation for ordinary differential equations as well, while the last two work only for delay equations. Unlike first order differential equations, where the oscillatory character is due to the delay only, the equation (2) can be oscillatory without any delay at all, i.e., in the case $\tau(t) \equiv t$. Figuratively speaking, two factors contribute to the oscillatory character of Eq.(2): the presence of the delay and the second order nature of the equation. The conditions (3.3), (3.4) and $(C_2)'$, $(C_3)'$ illustrate the role of these factors taken separately.

In what follows it will be assumed that the condition

$$\int^{+\infty} \tau(s)p(s)ds = +\infty \quad (3.5)$$

is fulfilled. As it follows from Lemma 4.1 in [24], this condition is necessary for Eq.(2) to be oscillatory. The study being devoted to the problem of oscillation of Eq.(2), the condition (3.5) does not affect the generality.

In this section oscillation results are obtained for Eq. (2) by reducing it to a first order equation. Since for the latter the oscillation is due solely to the delay, the criteria hold for delay equations only and do not work in the ordinary case.

Theorem 3.1 ([27]) Let (3.5) be fulfilled and the differential inequality

$$x'(t) + \left(\tau(t) + \int_T^{\tau(t)} \xi \tau(\xi)p(\xi)d\xi \right) p(t)x(\tau(t)) \leq 0$$

have no eventually positive solution. Then Eq. (2) is oscillatory.

Theorem 3.1 reduces the question of oscillation of Eq.(2) to that of the absence of eventually positive solutions of the differential inequality

$$x'(t) + \left(\tau(t) + \int_T^{\tau(t)} \xi \tau(\xi)p(\xi)d\xi \right) p(t)x(\tau(t)) \leq 0. \quad (3.6)$$

So oscillation results for first order delay differential equations can be applied since the oscillation of the equation

$$u'(t) + g(t)u(\delta(t)) = 0 \quad (3.7)$$

is equivalent to the absence of eventually positive solutions of the inequality

$$u'(t) + g(t)u(\delta(t)) \leq 0. \quad (3.8)$$

This fact is a simple consequence of the following comparison theorem deriving the oscillation of (3.7) from the oscillation of the equation

$$v'(t) + h(t)v(\sigma(t)) = 0. \quad (3.9)$$

We assume that $g, h : R_+ \rightarrow R_+$ are locally integrable, $\delta, \sigma : R_+ \rightarrow R$ are continuous, $\delta(t) \leq t$, $\sigma(t) \leq t$ for $t \in R_+$, and $\delta(t) \rightarrow +\infty$, $\sigma(t) \rightarrow +\infty$ as $t \rightarrow +\infty$.

Theorem 3.2 Let

$$g(t) \geq h(t) \quad \text{and} \quad \delta(t) \leq \sigma(t) \quad \text{for } t \in R_+$$

and let the equation (3.9) be oscillatory. Then (3.7) is also oscillatory.

Corollary 3.1 Let the equation (3.7) be oscillatory. Then the inequality (3.8) has no eventually positive solution.

Turning to applications of Theorem 3.1, we will use it together with the criteria (C_2) and (C_3) to get

Theorem 3.3 ([27]) Let

$$K := \limsup_{t \rightarrow \infty} \int_{\tau(t)}^t \left(\tau(s) + \int_0^{\tau(s)} \xi \tau(\xi) p(\xi) d\xi \right) p(s) ds > 1, \quad (C_2)''$$

or

$$k := \liminf_{t \rightarrow \infty} \int_{\tau(t)}^t \left(\tau(s) + \int_0^{\tau(s)} \xi \tau(\xi) p(\xi) d\xi \right) p(s) ds > \frac{1}{e}. \quad (C_3)''$$

Then Eq. (2) is oscillatory.

To apply Theorem (3.1) it suffices to note that: (i) (3.5) is fulfilled since otherwise $k = K = 0$; (ii) since $\tau(t) \rightarrow +\infty$ as $t \rightarrow +\infty$, the relations $(C_2)''$, $(C_3)''$ imply the same relations with 0 changed by any $T \geq 0$.

Remark 3.1 ([27]) Theorem 3.3 improves the criteria $(C_2)'$, $(C_3)'$ by Koplatadze [23] and Wei [59] mentioned above. This is directly seen from $(C_2)''$, $(C_3)''$ and can be easily checked if we take $\tau(t) \equiv t - \tau_0$ and $p(t) \equiv p_0 / (t - \tau_0)$ for $t \geq 2\tau_0$, where the constants $\tau_0 > 0$ and $p_0 > 0$ satisfy

$$\tau_0 p_0 < \frac{1}{e}.$$

In this case neither of $(C_2)'$, $(C_3)'$ is applicable for Eq. (2) while both $(C_2)''$, $(C_3)''$ give the positive conclusion about its oscillation. Note also that this is exactly the case where the oscillation is due to the delay since the corresponding equation without delay is nonoscillatory.

Remark 3.2 ([27]) The criteria $(C_2)''$, $(C_3)''$ look like (C_2) , (C_3) but there is an essential difference between them pointed out in the introduction. The condition (C_3) is close to the necessary one since according to [25] if $A \leq \frac{1}{e}$, then (3.7) is nonoscillatory. On the other hand, for an oscillatory equation (2) without

delay we have $k = K = 0$. Nevertheless, the constant $1/e$ in Theorem 3.3 is also the best possible in the sense that for any $\varepsilon \in (0, 1/e]$ it can not be replaced by $1/e - \varepsilon$ without affecting the validity of the theorem. This is illustrated by the following

Example 3.1 ([27]) Let $\varepsilon \in (0, 1/e]$, $1 - e\varepsilon < \beta < 1$, $\tau(t) \equiv \alpha t$ and $p(t) \equiv \beta(1 - \beta)\alpha^{-\beta}t^{-2}$, where $\alpha = e^{\frac{1}{\beta-1}}$. Then $(C_3)''$ is fulfilled with $1/e$ replaced by $1/e - \varepsilon$. Nevertheless Eq. (2) has a nonoscillatory solution, namely $u(t) \equiv t^\beta$. Indeed, denoting $c = \beta(1 - \beta)\alpha^{-\beta}$, we see that the expression under the limit sign in $(C_3)''$ is constant and equals $\alpha c |\ln \alpha| (1 + \alpha c) = (\beta/e)(1 + (\beta(1 - \beta))/e) > \beta/e > 1/e - \varepsilon$.

Note that there is a gap between the conditions $(C_2)'', (C_3)''$ when $0 \leq k \leq 1/e$, $k < K$. In the case of first order equations the conditions $(C_4) - (C_{13})$ attempt to fill this gap. Using results in this direction, one can derive various sufficient conditions for the oscillation of Eq. (2). According to Remark 3.1, neither of them can be optimal in the above sense but, nevertheless, they are of interest since they cannot be derived from other known results in the literature. We combine Theorem 3.1 with the result ([19], Corollary 1) to obtain

Theorem 3.4 ([27]) Let K and k be defined by $(C_2)'', (C_3)''$, $0 \leq k \leq 1/e$ and

$$K > k + \frac{1}{\lambda(k)} - \frac{1 - k - \sqrt{1 - 2k - k^2}}{2} \quad (C_{11})'$$

where $\lambda(k)$ is the smaller root of the equation $\lambda = \exp(k\lambda)$. Then Eq. (2) is oscillatory.

Note that the condition $(C_{11})'$ is analogous to the condition (C_{11}) .

REFERENCES

1. J. S. Bradley, Oscillation theorems for a second order equation. *J. Differential Equations*, **8** (1970), 397-403.
2. J. Diblík, Positive and oscillating solutions of differential equations with delay in critical case, *J. Comput. Appl. Math.* **88** (1998), 185-2002.
3. Y. Domshlak, Sturmian comparison method in oscillation study for discrete difference equations, I, *J. Diff. Integr. Eqs*, **7** (1994), 571-582.
4. Y. Domshlak, Delay-difference equations with periodic coefficients: sharp results in oscillation theory, *Math. Inequal. Appl.*, **1** (1998), 403-422.

5. Y. Domshlak and A. Aliev, On oscillatory properties of the first order differential equations with one or two retarded arguments, *Hiroshima Math. J.* **18** (1998), 31-46.
6. Y. Domshlak and I.P. Stavroulakis, Oscillations of first-order delay differential equations in a critical state, *Applicable Anal.*, **61** (1996), 359-377.
7. R. Edwards, Functional analysis. Theory and applications. Holt, Rinehart and Winston, Chicago - San Francisco - Toronto, 1965.
8. A. Elbert and I.P. Stavroulakis, Oscillations of first order differential equations with deviating arguments, Univ of Ioannina T.R. No 172 1990, *Recent trends in differential equations*, 163-178, World Sci. Ser. Appl. Anal.,1, World Sci. Publishing Co. (1992).
9. A. Elbert and I. P. Stavroulakis, Oscillation and non-oscillation criteria for delay differential equations, *Proc. Amer. Math. Soc.*, **123** (1995), 1503-1510.
10. L. Erbe, Oscillation criteria for second order nonlinear delay equations, *Canad. Math. Bull.* **16** (1973), 49-56.
11. L.H. Erbe, Qingkai Kong and B.G. Zhang, Oscillation Theory for Functional Differential Equations, Marcel Dekker, New York, 1995.
12. L.H. Erbe and B.G. Zhang, Oscillation of first order linear differential equations with deviating arguments, *Differential Integral Equations*, **1** (1988), 305-314.
13. N. Fukagai and T. Kusano, Oscillation theory of first order functional differential equations with deviating arguments, *Ann. Mat. Pura Appl.*, **136** (1984), 95-117.
14. K.Gopalsamy, Stability and Oscillations in Delay Differential Equations of Population Dynamics, Kluwer Academic Publishers, 1992.
15. I. Gyori and G. Ladas. Oscillation Theory of Delay Differential Equations with Applications, Clarendon Press, Oxford, 1991.
16. J.K. Hale, Theory of Functional Differential Equations, Springer-Verlag, New York, 1997.
17. E. Hille, Nonoscillation theorems, *Trans. Amer. Math. Soc.* **64** (1948), 234-252.
18. A.F. Ivanov, and V.N. Shevelo, Oscillation and asymptotic behavior of solutions of first order differential equations, *Ukrain, Math. Zh.*, **33** (1981), 745-751, 859.

19. J. Jaroš and I.P. Stavroulakis, Oscillation tests for delay equations, *Rocky Mountain J. Math.*, **29** (1999), 139-145.
20. C. Jian, Oscillation of linear differential equations with deviating argument, *Math. in Practice and Theory*, **1** (1991), 32-41 (in Chinese).
21. A. Kneser, Untersuchungen über die reellen Nulstellen der Integrale linearer Differentialgleichungen. *Math. Ann.* **42** (1893), 409-435.
22. M. Kon, Y.G. Sficas and I.P. Stavroulakis, Oscillation criteria for delay equations, *Proc. Amer. Math. Soc.*, **128** (2000), 2989-2997.
23. R. Koplatadze, Oscillation criteria of solutions of second order linear delay differential inequalities with a delayed argument (Russian). *Trudy Inst. Prikl. Mat. I.N.Vekua* **17** (1986), 104-120.
24. R. Koplatadze, On oscillatory properties of solutions of functional differential equations. *Mem. Differential Equations Math. Phys.* **3** (1994), 1-177.
25. R.G. Koplatadze and T.A. Chanturija, On the oscillatory and monotonic solutions of first order differential equations with deviating arguments, *Differentsial'nye Uravneniya*, **18** (1982), 1463-1465.
26. R.G. Koplatadze and G. Kvinikadze, On the oscillation of solutions of first order delay differential inequalities and equations, *Georgian Math. J.* **1** (1994), 675-685.
27. R.G. Koplatadze, G. Kvinikadze and I. P. Stavroulakis, Oscillation of second order linear delay differential equations, *Funct. Diff. Equ.* **7** (2000), 121-145.
28. E. Kozakiewicz, Conditions for the absence of positive solutions of first order differential inequalities with deviating arguments, 4th Intl. Coll. on Differential Equations, VSP, 1994, 157-161.
29. M.R. Kulenovic and M.K. Grammatikopoulos, First order functional differential inequalities with oscillating coefficients, *Nonlinear Anal.* **8** (1984), 1043-1054.
30. M.K. Kwong, Oscillation of first order delay equations, *J. Math. Anal. Appl.*, **156** (1991), 374-286.
31. M. K. Kwong and W. T. Patula, Comparison theorems for first order linear delay equations, *J. Differential Equations*, **70** (1987), 275-292.
32. G. Ladas, Sharp conditions for oscillations caused by delay, *Applicable Anal.*, **9** (1979), 93-98.

33. G. Ladas, V. Laskhmikantham and J.S. Papadakis, Oscillations of higher-order retarded differential equations generated by retarded arguments, *Delay and Functional Differential Equations and Their Applications*, Academic Press, New York, 1972, 219-231.
34. G. Ladas, C. Qian and J. Yan, A comparison result for the oscillation of delay differential equations, *Proc. Amer. Math. Soc.*, **114** (1992), 939-946.
35. G. Ladas, Y.G. Sficas and I.P. Stavroulakis, Functional-differential inequalities and equations with oscillating coefficients, *Trends in theory and practice of nonlinear differential equations*, (Arlington, Tex., 1982), 277-284, Lecture Notes in Pure and Appl. Math., 90, Marcel Dekker, New York, 1984.
36. G.S. Ladde, Oscillations caused by retarded perturbations of first order linear ordinary differential equations, *Atti Acad. Naz. Lincei. Rend. Cl. Sci. Fis. Mat. Natur.*, **63** (1977), 351-359.
37. G.S. Ladde, Class of functional equations with applications, *Nonlinear Anal.*, **2** (1978), 259-261.
38. G.S. Ladde, Stability and oscillation in single-species processes with past memory, *Int. J. System Sci.*, **10** (1979), 621-647.
39. G.S. Ladde, V. Lakshmikantham and B.G. Zhang, Oscillation Theory of Differential Equations with Deviating Arguments, Marcel Dekker, New York, 1987.
40. B. Li, Oscillations of delay differential equations with variable coefficients, *J. Math. Anal. Appl.*, **192** (1995), 312-321.
41. B. Li, Oscillations of first order delay differential equations, *Proc. Amer. Math. Soc.*, **124** (1996), 3729-3737.
42. A.D. Myshkis, Linear homogeneous differential equations of first order with deviating arguments, *Uspekhi Mat. Nauk*, **5** (1950), 160-162 (Russian).
43. A. D. Myshkis, Linear differential equations with retarded argument, Nauka, Moscow, 1972 (Russian).
44. S. B. Norkin, Differential equations of the second order with retarded arguments. Nauka, Moscow, 1965 (Russian).
45. Ch.G. Philos and Y.G. Sficas, An oscillation criterion for first-order linear delay differential equations, *Canad. Math. Bull.* **41** (1998), 207-213.
46. Y.G. Sficas and I.P. Stavroulakis, Oscillation criteria for first-order delay equations, *Bull. London Math. Soc.*, **35** (2003), 239-246.

47. C. Sturm, Sur les équations différentielles linéaires du second ordre, *J. Math. Pures Appl.* **1**(1836), 106-186.
48. C. A. Swanson, Comparison and oscillation theory of linear differential equations, Academic Press, New York and London, 1968.
49. X.H. Tang, Oscillation of delay differential equations with variable coefficients, *J. Math. Study*, **31** (3)(1998), 290-293.
50. X.H. Tang and J.S. Yu, Oscillations of first order delay differential equations in a critical state, *Mathematica Applicata* **13** (1)(2000), 75-79.
51. X.H. Tang and J.S. Yu, Oscillation of first order delay differential equations with oscillating coefficients, *Appl Math.-JCU*, **15** (2000), 252-258.
52. X.H. Tang and J.S. Yu, New oscillation criteria for delay difference equations, *Comput. Math. Applic.*, **42** (2001), 1319-1330.
53. X.H. Tang, J.S. Yu and Z.C. Wang, Comparison theorems of oscillation of first order delay differential equations in a critical state with applications, *Ke Xue Tongbao*, **44** (1999), 26-30.
54. A. Tomaras, Oscillation behavior of an equation arising from an industrial problem, *Bull. Austral. Math. Soc.*, **13** (1975), 255-260.
55. A. Tomaras, Oscillations of a first order functional differential equation, *Bull. Austral. Math. Soc.*, **17** (1977), 91-95.
56. A. Tomaras, Oscillatory behaviour of first order delay differential equations, *Bull. Austral. Math. Soc.*, **19** (1978), 183-190.
57. P. Waltman, A note on an oscillation criterion for an equation with a functional argument. *Canad. Math. Bull.* **11**(1968), 593-595
58. Z.C. Wang, I.P. Stavroulakis and X.Z. Qian, A survey on the oscillation of solutions of first order linear differential equations with deviating arguments, *Appl. Math. E-Notes*, **2** (2002), 171-191.
59. J. J. Wei, Oscillation of second order delay differential equation. *Ann. Differential Equations* **4** (1988), 473-478.
60. J. S. W. Wong, Second order oscillation with retarded arguments, in "Ordinary differential equations", pp. 581-596, Washington, 1971; Academic Press, New York and London, 1972.
61. J. Yan, Oscillatory property for second order linear differential equations, *J. Math. Anal. Appl.* **122** (1987), 380-384.

62. J.S. Yu and X.H. Tang, Comparison theorems in delay differential equations in a critical state and application, *Proc. London Math. Soc.*, **63** (2001), 188-204.
63. J.S. Yu and Z.C. Wang, Some further results on oscillation of neutral differential equations, *Bull. Austral. Math. Soc.*, **46** (1992), 149-157.
64. J.S. Yu, Z.C. Wang, B.G. Zhang and X.Z. Qian, Oscillations of differential equations with deviating arguments, *Panamerican Math. J.*, **2** (1992), 59-78.
65. Y. Zhou and Y.H. Yu, On the oscillation of solutions of first order differential equations with deviating arguments, *Acta Math. Appl. Sinica* **15**, no.3, (1999), 288-302.