

# Asymptotic behavior of the solutions of a class of rational difference equations

G. Papaschinopoulos, G. Stefanidou,  
Democritus University of Thrace  
School of Engineering  
67100 Xanthi  
Greece

## Abstract

In this paper we study the asymptotic behavior of the positive solutions of the rational difference equations

$$x_{n+1} = \frac{ax_{n-m(k+1)+1}}{\prod_{s=0}^k x_{n-m(s+1)+1} + 1}, \quad x_{n+1} = \frac{ax_{n-2k-1} \prod_{s=0}^k x_{n-2s}}{\prod_{s=0}^{2k+1} x_{n-s} + \prod_{s=0}^k x_{n-2s} + \prod_{s=0}^k x_{n-2s-1}},$$

$$x_{n+1} = \frac{ax_n x_{n-m(k+1)+1}}{x_n + x_{n-m(k+1)}}, \quad n = 0, 1, \dots,$$

where  $k, m \in \{1, 2, \dots\}$  and  $a$  is a positive number.

**Keywords:** Difference equation, periodic solution, convergence of the solutions.

## 1 Introduction

In [8] the author studied the global behavior of the second order rational difference equation having quadratic term

$$x_{n+1} = \frac{ax_{n-1}}{x_n x_{n-1} + b}, \quad a > 0, \quad b > 0 \quad (1.1)$$

and the third order difference equation having quadratic term

$$x_{n+1} = \frac{ax_n x_{n-1}}{x_n + bx_{n-2}}, \quad a > 0, \quad b > 0 \quad (1.2)$$

where  $x_{-2}, x_{-1}, x_0$  are real numbers. For the study of equation (1.1) the author used the fact that (1.1) reduces to a linear nonhomogeneous equation. Moreover, for the study of (1.2) he showed that equation (1.2) reduces to (1.1).

Furthermore in [3] the authors investigated equation (1.1) with nonnegative initial values  $x_{-1}, x_0$ . Moreover if we get  $b = 1$  in (1.1) then by dropping either the term  $x_n$  or  $x_{n-1}$  in the denominator of the equation (1.1) we obtain the equations

$$x_{n+1} = \frac{ax_{n-1}}{x_n + 1}, \quad x_{n+1} = \frac{ax_{n-1}}{x_{n-1} + 1}$$

which have been studied in [2]. Finally, results concerning rational difference equations having quadratic terms are included in [1], [3]-[11] and the references cited therein.

Now in this paper we study the following equations

$$x_{n+1} = \frac{ax_{n-m(k+1)+1}}{\prod_{s=0}^k x_{n-m(s+1)+1} + 1}, \quad n = 0, 1, \dots \quad (1.3)$$

$$x_{n+1} = \frac{ax_{n-2k-1} \prod_{s=0}^k x_{n-2s}}{\prod_{s=0}^{2k+1} x_{n-s} + \prod_{s=0}^k x_{n-2s} + \prod_{s=0}^k x_{n-2s-1}} \quad (1.4)$$

and

$$x_{n+1} = \frac{ax_n x_{n-m(k+1)+1}}{x_n + x_{n-m(k+1)}}, \quad n = 0, 1, \dots, \quad (1.5)$$

where  $a$  is a positive number,  $m, k \in \{1, 2, \dots\}$  and the initial values of the above equations are positive numbers. More precisely, we study the existence of periodic solutions and the asymptotic behavior of the positive solutions for equations (1.3)-(1.5). We note that equations (1.3)-(1.5) have a common property: They reduce to a linear nonhomogeneous equation.

## 2 Study of equation (1.3)

First we study the existence of positive periodic solutions of period  $m(k+1)$  for equation (1.3).

**Proposition 2.1** *Consider equation (1.3). Suppose that*

$$a > 1. \quad (2.1)$$

*Then equation (1.3) has periodic solutions of period  $m(k+1)$ .*

**Proof.** Let  $x_n$  be a positive solution of (1.3) with initial values  $x_{-m(k+1)+1}, x_{-m(k+1)+2}, \dots, x_0$  are positive numbers such that

$$\prod_{s=0}^k x_{i-m(s+1)+1} = a - 1, \quad i = 0, 1, \dots, m-1. \quad (2.2)$$

We prove that  $x_n$  is a periodic solution of (1.3) of period  $m(k+1)$ . From (1.3) and (2.2) we get

$$\begin{aligned} x_1 &= \frac{ax_{-m(k+1)+1}}{\prod_{s=0}^k x_{-m(s+1)+1} + 1} = x_{-m(k+1)+1}, \\ x_2 &= \frac{ax_{-m(k+1)+2}}{\prod_{s=0}^k x_{-m(s+1)+2} + 1} = x_{-m(k+1)+2}, \\ &\dots\dots\dots \\ x_m &= \frac{ax_{-mk}}{\prod_{s=0}^k x_{-ms} + 1} = x_{-mk}. \end{aligned} \quad (2.3)$$

Then from (1.3) and (2.3) we obtain

$$\begin{aligned} x_{m+1} &= \frac{ax_{-mk+1}}{x_1 \prod_{s=1}^k x_{-ms+1} + 1} = \frac{ax_{-mk+1}}{x_{-m(k+1)+1} \prod_{s=1}^k x_{-ms+1} + 1} = \\ &\frac{ax_{-mk+1}}{\prod_{s=0}^k x_{-m(s+1)+1} + 1} = x_{-mk+1}. \end{aligned}$$

Working inductively we can prove that

$$x_{m+j} = x_{-mk+j}, \quad j = 2, 3, \dots$$

and so the proof of the proposition is completed.

In the next proposition we study the asymptotic behavior of the positive solutions of (1.3). We need the following lemma.

**Lemma 2.1** *Let  $x_n$  be an arbitrary positive solution of (1.3). Then the following statements are true:*

(i) *If*

$$t_n = \prod_{s=0}^k x_{n-sm}^{-1}, \quad n = 1, 2, \dots \quad (2.4)$$

*with*

$$t_j = \prod_{s=0}^k x_{j-sm}^{-1}, \quad j = 1-m, 2-m, \dots, 0, \quad (2.5)$$

*then  $t_n$  satisfies the nonhomogeneous linear difference equation*

$$y_{n+1} = \frac{1}{a} y_{n+1-m} + \frac{1}{a}, \quad n = 0, 1, \dots \quad (2.6)$$

*Moreover,*

$$t_n = \begin{cases} B_n + \frac{n}{m}, & n = 1, 2, \dots \quad \text{if } a = 1 \\ \left(\frac{1}{a}\right)^{\frac{n}{m}} B_n + \frac{1}{a-1}, & n = 1, 2, \dots \quad \text{if } a \neq 1 \end{cases} \quad (2.7)$$

*where*

$$B_n = \sum_{i=0}^r c_i \cos\left(\frac{2\pi ni}{m}\right) + d_i \sin\left(\frac{2\pi ni}{m}\right), \quad r = \begin{cases} \frac{m-1}{2}, & \text{if } m \text{ is odd} \\ \frac{m}{2}, & \text{if } m \text{ is even} \end{cases} \quad (2.8)$$

*and  $c_i, d_i, i = 0, 1, \dots, r$  are constants which are derived from (2.5), (2.7) and (2.8).*

(ii) If

$$y_n^{(j)} = x_{m(k+1)n+j}, \quad j = 0, 1, \dots, m(k+1) - 1 \quad (2.9)$$

then

$$y_n^{(j)} = y_0^{(j)} \prod_{s=0}^{n-1} \frac{t_{m(k+1)(s+1)+j-m}}{t_{m(k+1)(s+1)+j}}, \quad j = 0, 1, \dots, m(k+1) - 1. \quad (2.10)$$

**Proof.** Let  $x_n$  be an arbitrary solution of (1.3). Then we get

$$x_{n+1} \prod_{s=1}^k x_{n+1-sm} = \frac{a x_{n-m(k+1)+1} \prod_{s=1}^k x_{n+1-sm}}{\prod_{s=0}^k x_{n+1-(s+1)m} + 1}$$

which implies that

$$\prod_{s=0}^k x_{n+1-sm} = \frac{a \prod_{s=0}^k x_{n+1-(s+1)m}}{\prod_{s=0}^k x_{n+1-(s+1)m} + 1}. \quad (2.11)$$

Then from (2.4) and (2.11) we have

$$\frac{1}{t_{n+1}} = \frac{\frac{a}{t_{n+1-m}}}{\frac{1}{t_{n+1-m}} + 1}$$

which implies that  $t_n$  satisfies the difference equation (2.6). Then relations (2.7) and (2.8) follow immediately. This completes the proof of statement (i).

(ii) From (2.4) we have

$$\frac{t_n}{t_{n-m}} = \frac{x_n^{-1} x_{n-m}^{-1} \cdots x_{n-km}^{-1}}{x_{n-m}^{-1} x_{n-2m}^{-1} \cdots x_{n-(k+1)m}^{-1}} = \frac{x_{n-m(k+1)}}{x_n}$$

which implies that

$$x_n = \frac{t_{n-m}}{t_n} x_{n-m(k+1)}, \quad n = 1, 2, \dots \quad (2.12)$$

So, from (2.9) and (2.12) it holds

$$y_{n+1}^{(j)} = \frac{t_{m(k+1)(n+1)+j-m}}{t_{m(k+1)(n+1)+j}} y_n^{(j)}, \quad j = 0, 1, \dots, m(k+1) - 1. \quad (2.13)$$

Therefore relation (2.13) implies that (2.10) is true. This completes the proof of the lemma.

**Proposition 2.2** *Consider equation (1.3). Then the following statements are true.*

(i) If

$$0 < a \leq 1 \quad (2.14)$$

then every positive solution of (1.3) tends to zero as  $n \rightarrow \infty$ .

(ii) If (2.1) holds then every positive solution of (1.3) tends to a periodic solution of period  $m(k+1)$ .

**Proof.** Let  $x_n$  be an arbitrary positive solution of (1.3).

(i) Suppose first that

$$0 < a < 1. \quad (2.15)$$

From (1.3) we get for  $j = 0, 1, \dots, m(k+1) - 1$

$$x_{m(k+1)n+j} < ax_{m(k+1)(n-1)+j} < \dots < a^n x_j. \quad (2.16)$$

Then from (2.15) and (2.16) we take

$$\lim_{n \rightarrow \infty} x_{m(k+1)n+j} = 0, \quad j = 0, 1, \dots, m(k+1) - 1$$

which implies that  $x_n$  tends to zero as  $n \rightarrow \infty$ .

Let now  $a = 1$ . We consider the functions

$$A_n^{(j)} = \ln \left( \prod_{s=0}^{n-1} \frac{t_{m(k+1)(s+1)+j-m}}{t_{m(k+1)(s+1)+j}} \right) = \sum_{s=0}^{n-1} \ln \left( \frac{t_{m(k+1)(s+1)+j-m}}{t_{m(k+1)(s+1)+j}} \right). \quad (2.17)$$

From (2.8) it is obvious that

$$B_{m(k+1)(s+1)+j-m} = B_{m(k+1)(s+1)+j}, \quad s = 0, 1, \dots, \quad j = 0, 1, \dots, m(k+1) - 1. \quad (2.18)$$

Hence relations (2.7), (2.8) and (2.18) imply that

$$t_{m(k+1)(s+1)+j-m} - t_{m(k+1)(s+1)+j} = -1, \quad s = 0, 1, \dots, \quad j = 0, 1, \dots, m(k+1) - 1. \quad (2.19)$$

In addition, if  $a$  is a real number such that  $1 + a > 0$  then

$$\ln(1 + a) < a. \quad (2.20)$$

Then from (2.19) and (2.20) we get

$$\begin{aligned} \sum_{s=0}^{n-1} \ln \left( 1 + \frac{t_{m(k+1)(s+1)+j-m} - t_{m(k+1)(s+1)+j}}{t_{m(k+1)(s+1)+j}} \right) &\leq \\ \sum_{s=0}^{n-1} \left( \frac{t_{m(k+1)(s+1)+j-m} - t_{m(k+1)(s+1)+j}}{t_{m(k+1)(s+1)+j}} \right) &= - \sum_{s=0}^{n-1} \frac{1}{t_{m(k+1)(s+1)+j}}. \end{aligned} \quad (2.21)$$

Since from (2.7)

$$\sum_{s=0}^{\infty} \frac{1}{t_{m(k+1)(s+1)+j}} = \infty$$

then from (2.21)

$$\sum_{s=0}^{\infty} \ln \left( \frac{t_{m(k+1)(s+1)+j-m}}{t_{m(k+1)(s+1)+j}} \right) = -\infty. \quad (2.22)$$

Therefore, from (2.17) and (2.22) we have

$$\lim_{n \rightarrow \infty} A_n^{(j)} = -\infty, \quad j = 0, 1, \dots, m(k+1) - 1$$

which implies that

$$\prod_{s=0}^{\infty} \frac{t_{m(k+1)(s+1)+j-m}}{t_{m(k+1)(s+1)+j}} = 0, \quad j = 0, 1, \dots, m(k+1) - 1. \quad (2.23)$$

So from (2.10) and (2.23) we have that  $x_n$  tends to zero as  $n \rightarrow \infty$ . This completes the proof of statement (i).

(ii) If  $a, b > 0$  then using (2.20) we can easily prove that

$$\left| \ln \left( \frac{a}{b} \right) \right| \leq |a - b| \max \left\{ \frac{1}{a}, \frac{1}{b} \right\}. \quad (2.24)$$

Then from (2.24) we have for  $j = 0, 1, \dots, m(k+1) - 1$

$$\begin{aligned} \left| \sum_{s=0}^{n-1} \ln \left( \frac{t_{m(k+1)(s+1)+j-m}}{t_{m(k+1)(s+1)+j}} \right) \right| &\leq \sum_{s=0}^{n-1} \left| \ln \left( \frac{t_{m(k+1)(s+1)+j-m}}{t_{m(k+1)(s+1)+j}} \right) \right| \leq \\ \sum_{s=0}^{n-1} |t_{m(k+1)(s+1)+j-m} - t_{m(k+1)(s+1)+j}| &\max \left\{ \frac{1}{t_{m(k+1)(s+1)+j}}, \frac{1}{t_{m(k+1)(s+1)+j-m}} \right\}. \end{aligned} \quad (2.25)$$

Furthermore, from (2.1), (2.7) and (2.18) we have

$$|t_{m(k+1)(s+1)+j-m} - t_{m(k+1)(s+1)+j}| = \left(\frac{1}{a}\right)^{\frac{m(k+1)(s+1)+j}{m}} |B_{m(k+1)(s+1)+j}|(a-1). \quad (2.26)$$

Then using (2.7) and (2.26) we can prove that there exists a positive number  $M > 0$  such that

$$|t_{m(k+1)(s+1)+j-m} - t_{m(k+1)(s+1)+j}| \max \left\{ \frac{1}{t_{m(k+1)(s+1)+j}}, \frac{1}{t_{m(k+1)(s+1)+j-m}} \right\} \leq M \left(\frac{1}{a}\right)^{\frac{m(k+1)(s+1)+j}{m}}, \quad j = 0, 1, \dots, m(k+1) - 1. \quad (2.27)$$

Therefore, from (2.1), (2.25) and (2.27) it follows that

$$\left| \sum_{s=0}^{\infty} \ln \left( \frac{t_{m(k+1)(s+1)+j-m}}{t_{m(k+1)(s+1)+j}} \right) \right| < \infty. \quad (2.28)$$

Then using (2.17) and (2.28) it is obvious that there exist the

$$\lim_{n \rightarrow \infty} A_n^{(j)} = l_j < \infty, \quad j = 0, 1, \dots, m(k+1) - 1. \quad (2.29)$$

Relations (2.9), (2.10), (2.17) and (2.29) imply that

$$\lim_{n \rightarrow \infty} x_{m(k+1)n+j} = p_j < \infty, \quad j = 0, 1, \dots, m(k+1) - 1.$$

This completes the proof of the proposition.

### 3 Study of equation (1.4)

First we study the existence of positive solutions of period  $2k+2$  for the equation (1.4).

**Proposition 3.1** *Consider equation (1.4) where*

$$a > 2. \quad (3.1)$$

*Then equation (1.4) has positive periodic solutions of period  $2k+2$ .*



**Proof.** Let  $x_n$  be a positive solution of (1.4) with initial values such that

$$\prod_{s=0}^k x_{-2s} = \prod_{s=0}^k x_{-2s-1} = a - 2. \quad (3.2)$$

Then from (1.4) and (3.2) we get

$$\begin{aligned} x_1 &= \frac{a x_{-2k-1} \prod_{s=0}^k x_{-2s}}{\prod_{s=0}^{2k+1} x_{-s} + \prod_{s=0}^k x_{-2s} + \prod_{s=0}^k x_{-2s-1}} = \frac{a(a-2)x_{-2k-1}}{(a-2)^2 + 2(a-2)} = x_{-2k-1}, \\ x_2 &= \frac{a x_1 x_{-2k} \prod_{s=1}^k x_{1-2s}}{x_1 \prod_{s=1}^{2k+1} x_{1-s} + x_1 \prod_{s=1}^k x_{1-2s} + \prod_{s=0}^k x_{-2s}} = \\ &= \frac{a x_{-2k-1} x_{-2k} \prod_{s=1}^k x_{1-2s}}{x_{-2k-1} \prod_{s=1}^{2k+1} x_{1-s} + x_{-2k-1} \prod_{s=1}^k x_{1-2s} + \prod_{s=0}^k x_{-2s}} = \\ &= \frac{a x_{-2k} \prod_{s=0}^k x_{-2s-1}}{\prod_{s=0}^{2k+1} x_{-s} + \prod_{s=0}^k x_{-2s-1} + \prod_{s=0}^k x_{-2s}} = \frac{a(a-2)x_{-2k}}{(a-2)^2 + 2(a-2)} = x_{-2k}. \end{aligned}$$

Working inductively we can prove that

$$x_n = x_{n-2k-2}, \quad n = 3, 4, \dots$$

This completes the proof of the proposition.

In the following proposition we study the asymptotic behavior of the positive solutions of (1.4). We need the following lemma.

**Lemma 3.1** *Let  $x_n$  be a positive solution of (1.4). Then the following statements are true:*

(i) If

$$t_n = \prod_{s=0}^k x_{n-2s}^{-1}, \quad n = 1, 2, \dots \quad (3.3)$$

with

$$t_j = \prod_{s=0}^k x_{j-2s}^{-1}, \quad j = -1, 0, \quad (3.4)$$

then  $t_n$ ,  $n = 1, 2, \dots$  satisfies the following difference equation

$$y_{n+1} = \frac{1}{a}y_n + \frac{1}{a}y_{n-1} + \frac{1}{a}, \quad n = 0, 1, \dots \quad (3.5)$$

Moreover,

$$t_n = \begin{cases} c_1(-\frac{1}{2})^n + c_2 + \frac{1}{3}n, & n = 1, 2, \dots \quad \text{if } a = 2 \\ c_1p_1^n + c_2p_2^n + \frac{1}{a-2}, & n = 1, 2, \dots \quad \text{if } a \neq 2 \end{cases} \quad (3.6)$$

where

$$p_1 = \frac{1}{2a}(1 - \sqrt{1+4a}), \quad p_2 = \frac{1}{2a}(1 + \sqrt{1+4a}), \quad (3.7)$$

$c_1, c_2$  are defined from (3.4) and (3.6).

(ii) If

$$y_n^{(j)} = x_{2(k+1)n+j}, \quad j = 0, 1, \dots, 2k+1 \quad (3.8)$$

then

$$y_n^{(j)} = y_0^{(j)} \prod_{s=0}^{n-1} \frac{t_{2(k+1)(s+1)+j-2}}{t_{2(k+1)(s+1)+j}}, \quad j = 0, 1, \dots, 2k+1. \quad (3.9)$$

**Proof.** (i) Let  $x_n$  be an arbitrary positive solution of (1.4). Then we get

$$x_{n+1} \prod_{s=1}^k x_{n-2s+1} = \frac{ax_{n-2k-1} \prod_{s=0}^k x_{n-2s} \prod_{s=1}^k x_{n-2s+1}}{\prod_{s=0}^{2k+1} x_{n-s} + \prod_{s=0}^k x_{n-2s} + \prod_{s=0}^k x_{n-2s-1}}$$

which implies that

$$\prod_{s=0}^k x_{n-2s+1} = \frac{a \prod_{s=0}^{2k+1} x_{n-s}}{\prod_{s=0}^{2k+1} x_{n-s} + \prod_{s=0}^k x_{n-2s} + \prod_{s=0}^k x_{n-2s-1}}. \quad (3.10)$$

Then relations (3.3) and (3.10) imply that

$$\frac{1}{t_{n+1}} = \frac{\frac{a}{t_n t_{n-1}}}{\frac{1}{t_n t_{n-1}} + \frac{1}{t_n} + \frac{1}{t_{n-1}}}$$

from which we take that  $t_n$  satisfies the difference equation (3.5). Then relation (3.6) follows immediately.

(ii) Using (3.3) we take

$$\frac{t_n}{t_{n-2}} = \frac{x_n^{-1} x_{n-2}^{-1} \cdots x_{n-2k}^{-1}}{x_{n-2}^{-1} x_{n-4}^{-1} \cdots x_{n-2k-2}^{-1}} = \frac{x_{n-2k-2}}{x_n}$$

which implies that

$$x_n = \frac{t_{n-2}}{t_n} x_{n-2k-2}, \quad n = 1, 2, \dots \quad (3.11)$$

So, from (3.8) and (3.11) it holds

$$y_n^{(j)} = \frac{t_{2(k+1)n+j-2}}{t_{2(k+1)n+j}} y_{n-1}^{(j)}, \quad j = 0, 1, \dots, 2k+1. \quad (3.12)$$

From (3.12) relation (3.9) follows immediately. This completes the proof of the lemma.

**Proposition 3.2** *Consider equation (1.4). Then the following statements are true:*

(ii) *If*

$$0 < a \leq 2 \quad (3.13)$$

*then every positive solution of (1.4) tends to zero as  $n \rightarrow \infty$ .*

(ii) If

$$a > 2 \quad (3.14)$$

then every positive solution of (1.4) tends to a periodic solution of (1.4) of period  $2k + 2$ .

**Proof.** Let  $x_n$  be an arbitrary positive solution of (1.4).

(i) Suppose that (2.15) is satisfied. Relation (1.4) implies that for  $j = 0, 1, \dots, 2k + 1$

$$x_{2(k+1)n+j} < ax_{2(k+1)(n-1)+j} < \dots < a^n x_j. \quad (3.15)$$

Therefore, from (2.15) and (3.15) we get

$$\lim_{n \rightarrow \infty} x_{2(k+1)n+j} = 0, \quad j = 0, 1, \dots, 2k + 1 \quad (3.16)$$

which imply that  $x_n$  tends to zero as  $n \rightarrow \infty$ .

Suppose that

$$1 \leq a < 2. \quad (3.17)$$

From (3.7) and (3.17) we can easily prove that

$$|p_1| < 1, \quad 1 < p_2. \quad (3.18)$$

We set for  $j = 0, 1, \dots, 2k + 1$

$$B_n^{(j)} = \ln \left( \prod_{s=0}^{n-1} \frac{t_{2(k+1)(s+1)+j-2}}{t_{2(k+1)(s+1)+j}} \right). \quad (3.19)$$

Then from (2.20) we have for  $j = 0, 1, \dots, 2k + 1$

$$\begin{aligned} B_n^{(j)} &= \sum_{s=0}^{n-1} \ln \left( 1 + \frac{t_{2(k+1)(s+1)+j-2} - t_{2(k+1)(s+1)+j}}{t_{2(k+1)(s+1)+j}} \right) \leq \\ &\sum_{s=0}^{n-1} \left( \frac{t_{2(k+1)(s+1)+j-2} - t_{2(k+1)(s+1)+j}}{t_{2(k+1)(s+1)+j}} \right). \end{aligned} \quad (3.20)$$

Moreover, from (3.6) and (3.20) we can prove that

$$\frac{t_{2(k+1)(s+1)+j-2} - t_{2(k+1)(s+1)+j}}{t_{2(k+1)(s+1)+j}} = \frac{c_1(p_1^{-2} - 1) \left( \frac{p_1}{p_2} \right)^{2(k+1)(s+1)+j} + c_2(p_2^{-2} - 1)}{c_1 \left( \frac{p_1}{p_2} \right)^{2(k+1)(s+1)+j} + c_2 + \frac{1}{a-2} p_2^{-2(k+1)(s+1)+j}}. \quad (3.21)$$

Using (3.18) and (3.21) we have that

$$\lim_{s \rightarrow \infty} \left( \frac{t_{2(k+1)(s+1)+j-2} - t_{2(k+1)(s+1)+j}}{t_{2(k+1)(s+1)+j}} \right) = p_2^{-2} - 1 < 0. \quad (3.22)$$

Therefore, from (3.20) and (3.22) we can prove that

$$\lim_{n \rightarrow \infty} B_n^{(j)} = -\infty, \quad j = 0, 1, \dots, 2k+1 \quad (3.23)$$

which from (3.19) imply that for  $j = 0, 1, \dots, 2k+1$

$$\prod_{s=0}^{\infty} \frac{t_{2(k+1)(s+1)+j-2}}{t_{2(k+1)(s+1)+j}} = 0. \quad (3.24)$$

Hence, from (3.8), (3.9) and (3.24) we have that relations (3.16) are true and so  $x_n$  tends to zero as  $n \rightarrow \infty$ .

Suppose now that

$$a = 2. \quad (3.25)$$

So from (3.6) and (3.25) we get

$$\begin{aligned} & \frac{t_{2(k+1)(s+1)+j-2} - t_{2(k+1)(s+1)+j}}{t_{2(k+1)(s+1)+j}} = \\ & \frac{3c_1 \left(-\frac{1}{2}\right)^{2(k+1)(s+1)+j} - \frac{2}{3}}{c_2 + c_1 \left(-\frac{1}{2}\right)^{2(k+1)(s+1)+j} + \frac{1}{3}(2(k+1)(s+1)+j)}. \end{aligned} \quad (3.26)$$

Then from (3.26) we can easily prove

$$\sum_{s=0}^{\infty} \left( \frac{t_{2(k+1)(s+1)+j-2} - t_{2(k+1)(s+1)+j}}{t_{2(k+1)(s+1)+j}} \right) = -\infty. \quad (3.27)$$

Therefore, from (3.20), (3.27) we have that (3.23) is satisfied and so arguing as above (3.16) hold which implies that  $x_n$  tends to zero as  $n \rightarrow \infty$ .

(ii) Finally, suppose that (3.14) is satisfied. Then from (3.7) it is obvious that

$$\left| \frac{p_1}{p_2} \right| < 1, \quad |p_1| < 1, \quad p_2 < 1. \quad (3.28)$$

In addition, from (3.6) we have that for  $j = 0, 1, \dots, 2k+1$

$$\begin{aligned} & t_{2(k+1)(s+1)+j-2} - t_{2(k+1)(s+1)+j} = \\ & p_2^{2(k+1)(s+1)+j} \left( c_1(p_1^{-2} - 1) \left( \frac{p_1}{p_2} \right)^{2(k+1)(s+1)+j} + c_2(p_2^{-2} - 1) \right). \end{aligned} \quad (3.29)$$

In addition, from (2.24) we get for  $j = 0, 1, \dots, 2k + 1$

$$\left| \ln \left( \frac{t_{2(k+1)(s+1)+j-2}}{t_{2(k+1)(s+1)+j}} \right) \right| \leq |t_{2(k+1)(s+1)-2+j} - t_{2(k+1)(s+1)+j}| \max \left\{ \frac{1}{t_{2(k+1)(s+1)-2+j}}, \frac{1}{t_{2(k+1)(s+1)+j}} \right\}. \quad (3.30)$$

Using (3.6), (3.28), (3.29) and (3.30) there exists a positive number  $N$  such that for  $j = 0, 1, \dots, 2k + 1$

$$\left| \ln \left( \frac{t_{2(k+1)(s+1)+j-2}}{t_{2(k+1)(s+1)+j}} \right) \right| \leq N p_2^{2(k+1)(s+1)+j}. \quad (3.31)$$

Therefore, from (3.19) and (3.31) we have that there exist

$$\lim_{n \rightarrow \infty} B_n^{(j)} = \mu_j < \infty, \quad j = 0, 1, \dots, 2k + 1. \quad (3.32)$$

Hence, relations (3.8), (3.9), (3.19) and (3.32) imply that

$$\lim_{n \rightarrow \infty} x_{2(k+1)n+j} = q_j < \infty, \quad j = 0, 1, \dots, 2k + 1$$

and so the proof of the proposition is completed.

## 4 Study of equation (1.5)

In the first proposition we study the existence of positive periodic solutions of (1.5) of period  $m(k + 1)$ .

**Proposition 4.1** *Consider equation (1.5) where (3.25) holds. Let  $x_n$  be positive solution of (1.5) such that*

$$x_0 = x_{-m(k+1)}. \quad (4.1)$$

*Then  $x_n$  is a periodic solution of (1.5) with period  $m(k + 1)$ .*

**Proof** Let  $x_n$  be a positive solution of (1.5) such that (4.1) holds. Then from (1.5), (3.25) we get

$$x_1 = \frac{2x_0 x_{-m(k+1)+1}}{x_0 + x_{-m(k+1)}} = \frac{2x_0 x_{-m(k+1)+1}}{2x_0} = x_{-m(k+1)+1}$$

and working inductively we can prove that

$$x_n = x_{n-m(k+1)}, \quad n = 1, 2, \dots$$

This completes the proof of the proposition.

In the last proposition of this paper we study the asymptotic behavior of the positive solutions of (1.5). We need the following lemma.

**Lemma 4.1** *Consider equation (1.5). Let  $x_n$  be a positive solution of (1.5). Then if  $a \neq 1$ , for  $j = 0, 1, \dots, m(k+1) - 1$  and  $n = 0, 1, \dots$  it holds*

$$x_{nm(k+1)+j} = (a-1)^n x_j \prod_{s=1}^n \frac{1}{c(a-1)\left(\frac{1}{a}\right)^{sm(k+1)+j} + 1} \quad (4.2)$$

where

$$c = \frac{x_{-m(k+1)}}{x_0} - \frac{1}{a-1}$$

and if  $a = 1$ , for  $j = 0, 1, \dots, m(k+1) - 1$  and  $n = 0, 1, \dots$  it holds

$$x_{nm(k+1)+j} = x_j \prod_{s=1}^n \frac{1}{d + sm(k+1) + j}, \quad d = \frac{x_{-m(k+1)}}{x_0}. \quad (4.3)$$

**Proof.** We set

$$y_n = \frac{x_{n-m(k+1)}}{x_n}. \quad (4.4)$$

Then from (1.5) and (4.4) we get

$$y_{n+1} = \frac{1}{a} y_n + \frac{1}{a}, \quad n = 0, 1, \dots \quad (4.5)$$

So from (4.4) and (4.5) relations (4.2) and (4.3) follow immediately. This completes the proof of the lemma.

**Proposition 4.2** *Consider equation (1.5). Then the following statements are true:*

- (i) *If  $0 < a < 2$  then every positive solution of (1.5) tends to zero as  $n \rightarrow \infty$ .*
- (ii) *If  $a = 2$  then every positive solution of (1.5) tends to a periodic solution of (1.5) of period  $m(k+1)$  as  $n \rightarrow \infty$ .*
- (iii) *If  $a > 2$  then every positive solution of (1.5) tends to  $\infty$  as  $n \rightarrow \infty$ .*

**Proof.** Let  $x_n$  be an arbitrary solution of (1.5).

(i) Suppose that (2.15) holds. Then using (1.5) and arguing as in Proposition 2.2 we can prove that  $x_n$  tends to zero as  $n \rightarrow \infty$ .

Suppose that

$$1 < a < 2. \quad (4.6)$$

Let for  $j = 0, 1, \dots, m(k+1) - 1$

$$D_n^{(j)} = \prod_{s=1}^n \frac{1}{c(a-1)\left(\frac{1}{a}\right)^{sm(k+1)+j} + 1}. \quad (4.7)$$

We have for  $j = 0, 1, \dots, m(k+1) - 1$

$$\ln(D_n^{(j)}) = -\sum_{s=1}^n \ln\left(c(a-1)\left(\frac{1}{a}\right)^{sm(k+1)+j} + 1\right). \quad (4.8)$$

In addition, from (2.20) we take

$$|\ln(1+a)| \leq \max\left\{a, \frac{-a}{1+a}\right\}. \quad (4.9)$$

Using (4.8) and (4.9) and since

$$1 < a \quad (4.10)$$

we can prove that

$$\lim_{n \rightarrow \infty} (\ln(D_n^{(j)})) = L_j < \infty, \quad j = 0, 1, \dots, m(k+1) - 1 \quad (4.11)$$

which implies that

$$\lim_{n \rightarrow \infty} D_n^{(j)} = M_j < \infty, \quad j = 0, 1, \dots, m(k+1) - 1. \quad (4.12)$$

Therefore, from (4.2), (4.6), (4.7) and (4.12) we have that

$$\lim_{n \rightarrow \infty} x_{nm(k+1)+j} = 0, \quad j = 0, 1, \dots, m(k+1) - 1 \quad (4.13)$$

and so  $x_n$  tends to zero as  $n \rightarrow \infty$ .

Let now  $a = 1$ . We set for  $j = 0, 1, \dots, m(k+1) - 1$

$$K_n^{(j)} = \prod_{s=1}^n \frac{1}{d + sm(k+1) + j}. \quad (4.14)$$



Then from (4.14) for  $j = 0, 1, \dots, m(k+1) - 1$  we take

$$\ln(K_n^{(j)}) = -\sum_{s=1}^n \ln(d + sm(k+1) + j). \quad (4.15)$$

So from (4.15) we can prove that

$$\lim_{n \rightarrow \infty} (\ln(K_n^{(j)})) = -\infty, \quad j = 0, 1, \dots, m(k+1) - 1$$

which implies that

$$\lim_{n \rightarrow \infty} K_n^{(j)} = 0, \quad j = 0, 1, \dots, m(k+1) - 1. \quad (4.16)$$

Then relations (4.3), (4.14), (4.16) imply that (4.13) are true and so  $x_n$  tends to zero as  $n \rightarrow \infty$ .

(ii) Suppose now that  $a = 2$ . Then from (4.10) relations (4.12) are true. So from (4.2) we have

$$\lim_{n \rightarrow \infty} x_{nm(k+1)+j} = M_j x_j < \infty, \quad j = 0, 1, \dots, m(k+1) - 1$$

and so  $x_n$  tends to a periodic solution of (1.5) of period  $m(k+1)$  as  $n \rightarrow \infty$ .

(iii) Finally, suppose that  $a > 2$ . Then using (4.10), we have that relations (4.12) hold and so from (4.2) it is clear that  $x_n$  tends to  $\infty$  as  $n \rightarrow \infty$ . This completes the proof of the proposition.

## References

- [1] A.M. Amleh, N. Kruse and G. Ladas, On a class of difference equations with strong negative feedback, J. Differ. Equ. Appl. 5 (1999), 497-515.
- [2] A.M. Amleh, E. Camouzis and G. Ladas, On second order rational difference equations, part I, J. Differ. Equ. Appl. 13 (2007), 969-1004.
- [3] A.M. Amleh, E. Camouzis and G. Ladas, On the dynamics of a rational difference equation, part I, Int. J. Differ. Equ. 3 (2008), 1-35.
- [4] K. S. Berenhaut, J.D. Foley, S. Stević, The global attractivity of the rational difference equation  $y_n = \frac{y_{n-k} + y_{n-m}}{1 + y_{n-k} y_{n-m}}$ . Appl. Math. Lett. 20 (2007), no. 1, 54-58.
- [5] E. Camouzis, G. Ladas, I.W. Rodrigues and S. Northshield, On the rational recursive sequence  $x_{n+1} = (\beta x_n^2)/(1 + x_{n-1}^2)$ , Comput. Math. Appl. 28 (1994), pp. 37-43.

- [6] M. Dehghan, C.M. Kent, R. Mazrooei-Sebdani, N. L. Ortiz, H. Sedaghat, Dynamics of rational difference equations containing quadratic terms, *J. Differ. Equ. Appl.*, 14 (2008), 191-208.
- [7] E.A. Grove, E.J. Janowski, C.M. Kent, G. Ladas, On the rational recursive sequence  $x_{n+1} = \frac{\alpha x_n + \beta}{(\gamma x_n + \delta)x_{n-1}}$ . *Comm. Appl. Nonlinear Anal.* 1 (1994), no. 3, pp. 61-72.
- [8] H. Sedaghat, Global behaviours of rational difference equations of orders two and three with quadratic terms, *J. Difference Equ. Appl.*, 15(3) (2009), 215-224.
- [9] S. Stević, Global stability and asymptotics of some classes of rational difference equations, *J. Math. Anal. Appl.* 316 (2006) 60-68.
- [10] X. Yang, On the global asymptotic stability of the difference equation  $x_n = \frac{x_{n-1}x_{n-2} + x_{n-3} + \alpha}{x_{n-1} + x_{n-2}x_{n-3} + \alpha}$ , *Appl. Math. Comput.* 171 (2005) 857-861.
- [11] X. Yang, D.J. Evans and G.M. Megson, On two rational difference equations, *Appl. Math. Comput.* 176 (2006) 422-430.